

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 October 2001 (25.10.2001)

PCT

(10) International Publication Number
WO 01/78740 A1

- (51) International Patent Classification⁷: **A61K 31/575**, 31/137, A61P 11/06 // (A61K 31/575, 31:137)
- (21) International Application Number: PCT/GB01/01637
- (22) International Filing Date: 11 April 2001 (11.04.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
0009609.9 18 April 2000 (18.04.2000) GB
- (71) Applicant (*for all designated States except US*): **GLAXO GROUP LIMITED** [GB/GB]; Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 ONN (GB).
- (74) Agent: **LEAROYD, Stephanie, Anne**; GlaxoSmithKline, Corporate Intellectual Property, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **GAVIN, Brian**, Charles [IE/IE]; GlaxoSmithKline, Grange Road, P.O. Box 700, Rathfarnham, 16 Dublin (IE). **GARRETT, Ronique, Nichele** [US/US]; GlaxoSmithKline, Five Moore Drive, Research Triangle Park, Durham County, NC 27709 (US). **ROCHE, Trevor, Charles** [GB/GB]; GlaxoSmithKline, Park Road, Ware, Hertfordshire SG12 0DP (GB).
- Published:**
— *with international search report*
— *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: MEDICAL COMBINATIONS COMPRISING MOMETASONE AND SALMETEROL

(57) Abstract: The present invention is concerned with pharmaceutical formulations comprising a combination of salmeterol and mometasone and the use of such formulations in medicine, particularly in the prophylaxis and treatment of respiratory diseases.



WO 01/78740 A1

MEDICAL COMBINATIONS COMPRISING MOMETASONE AND SALMETEROL

The present invention is concerned with combinations of salmeterol and mometasone, particularly compositions containing a combination of salmeterol and mometasone and the use of such compositions in medicine, particularly in the prophylaxis and treatment of respiratory diseases.

GB 2 140 800 describes phenethanolamine compounds which are β_2 -adrenoreceptor agonists including 4-hydroxy- α^1 -[[[6-(4-phenylbutoxy)hexyl]-amino]methyl]-1,3-benzenedimethanol 1-hydroxy-2-naphthalenecarboxylate (salmeterol xinafoate) which is now used clinically in the treatment of bronchial asthma and related disorders.

EP 57,401 and US 4,472,393 describe mometasone i.e. 9,21-dichloro-11 β ,17-dihydroxy-16 α -methylpregna-1,4-diene-3,20-dione, esters thereof such as mometasone furoate i.e. (11 β ,16 α)-9,21-dichloro-17-[(2-furanylcarbonyl)oxy]-11-hydroxy-16-methylpregna-1,4-diene-3,20-dione, and pharmaceutical formulations thereof. Mometasone is an antiinflammatory corticosteroid, which is now used clinically in the treatment of respiratory disorders.

Although salmeterol xinafoate and mometasone furoate are effective therapies, there exists a clinical need for asthma therapies having potent and selective action and having an advantageous profile of action.

Therefore, according to the present invention there is provided a combination of salmeterol or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof and mometasone or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof.

It will be appreciated that the compounds of the combination may be administered simultaneously, either in the same or different pharmaceutical formulations or sequentially. If there is sequential administration, the delay in administering the second compound should not be such as to lose the beneficial therapeutic effect of the combination.

According to a further aspect of the present invention, there is provided a pharmaceutical formulation comprising salmeterol or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof and mometasone or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, and a pharmaceutically acceptable carrier or excipient, and optionally one or more other therapeutic ingredients. According to a preferred aspect of the present invention, there is provided a pharmaceutical formulation comprising salmeterol xinafoate and mometasone furoate (suitably as in the form of the monohydrate), and a pharmaceutically acceptable carrier or excipient, and optionally one or more other therapeutic ingredients. In the most preferred aspect, the above pharmaceutical formulations are suitable for administration by inhalation.

It is to be understood that the present invention covers all combinations of particular and preferred aspects of the invention described herein.

As would be appreciated by the skilled person, salmeterol includes an asymmetric centre, and mometasone contains several asymmetric centres. The present invention includes both (S) and (R) enantiomers of salmeterol either in substantially pure form or admixed in any proportions, as well as each isomer of mometasone either in substantially pure form or admixed in any proportions. The enantiomers of salmeterol have been described previously, for example, in EP0422889 and WO 99/13867.

By the term "physiologically functional derivative" is meant a chemical derivative of salmeterol or mometasone having the same physiological function as the free compound, for example, by being convertible in the body thereto. According to the present invention, examples of physiologically functional derivatives include esters.

Suitable salts according to the invention include those formed with both organic and inorganic acids. Pharmaceutically acceptable acid addition salts include but are not limited to those formed from hydrochloric, hydrobromic, sulphuric, citric, tartaric, phosphoric, lactic, pyruvic, acetic, trifluoroacetic, succinic, oxalic, fumaric, maleic, oxaloacetic, methanesulphonic, ethanesulphonic, p-

toluenesulphonic, benzenesulphonic, isethionic, and naphthalenecarboxylic, such as 1-hydroxy-2-naphthalenecarboxylic acids.

5 Pharmaceutically acceptable esters of salmeterol or mometasone may have a hydroxyl group converted to a C₁₋₆alkyl, aryl, aryl C₁₋₆ alkyl, hetaryl (such as furanyl) or amino acid ester.

10 As mentioned above, both salmeterol and mometasone and their pharmaceutically acceptable salts, solvates, and physiologically functional derivatives have been described for use in the treatment of respiratory diseases. Therefore, formulations of salmeterol and mometasone and their pharmaceutically acceptable salts, solvates, and physiologically functional derivatives have use in the prophylaxis and treatment of clinical conditions for which a selective β_2 -adrenoreceptor agonist and/or an antiinflammatory
15 corticosteroid is indicated. Such conditions include diseases associated with reversible airways obstruction such as asthma, chronic obstructive pulmonary diseases (COPD) (e.g. chronic and wheezy bronchitis, emphysema), respiratory tract infection and upper respiratory tract disease.

20 Accordingly, the present invention provides a method for the prophylaxis or treatment of a clinical condition in a mammal, such as a human, for which a selective β_2 -adrenoreceptor agonist and/or antiinflammatory corticosteroid is indicated, which comprises administration of a therapeutically effective amount of a combination of salmeterol or a pharmaceutically acceptable salt, solvate, or
25 physiologically functional derivative thereof and mometasone or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof. The present invention further provides a method for the prophylaxis or treatment of a clinical condition in a mammal, such as a human, for which a selective β_2 -adrenoreceptor agonist and/or antiinflammatory corticosteroid is
30 indicated, which comprises administration of a therapeutically effective amount of a pharmaceutical formulation comprising salmeterol or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof and mometasone or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, and a pharmaceutically acceptable carrier or
35 excipient. In a preferred aspect, there is provided such a method which

comprises administration of a therapeutically effective amount of a pharmaceutical formulation comprising salmeterol xinafoate and mometasone furoate (suitably as the monohydrate), and a pharmaceutically acceptable carrier or excipient. In particular, the present invention provides such methods for the prophylaxis or treatment of a disease associated with reversible airways obstruction such as asthma, chronic obstructive pulmonary disease (COPD), respiratory tract infection or upper respiratory tract disease.

In the alternative, there is provided a combination of salmeterol or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof and mometasone or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, for use in therapy, particularly for use in the prophylaxis or treatment of a clinical condition for which a selective β_2 -adrenoreceptor agonist and/or antiinflammatory corticosteroid is indicated. In particular, there is provided a pharmaceutical formulation comprising salmeterol or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof (suitably, salmeterol xinafoate) and mometasone or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof (suitably, mometasone furoate optionally in the form of the monohydrate), and a pharmaceutically acceptable carrier or excipient for use in therapy, particularly for use in the prophylaxis or treatment of a clinical condition for which a selective β_2 -adrenoreceptor agonist and/or antiinflammatory corticosteroid is indicated. In a preferred aspect, the invention is concerned with the prophylaxis or treatment of a disease associated with reversible airways obstruction such as asthma, chronic obstructive pulmonary disease (COPD), respiratory tract infection or upper respiratory tract disease.

The amount of salmeterol and mometasone, or a pharmaceutically acceptable salt, solvate or physiologically functional derivative thereof which is required to achieve a therapeutic effect will, of course, vary with the particular compound, the route of administration, the subject under treatment, and the particular disorder or disease being treated. As a monotherapy, salmeterol xinafoate is generally administered to adult humans by aerosol inhalation at a dose of 50mcg or 100mcg twice daily.

While it is possible for the active ingredients of the combination to be administered as the raw chemical, it is preferable to present them as a pharmaceutical formulation. When the individual compounds of the combination are administered separately, they are generally each presented as a pharmaceutical formulation as described previously in the art.

Pharmaceutical formulations are often prescribed to the patient in "patient packs" containing the whole course of treatment in a single package. Patient packs have an advantage over traditional prescriptions, where a pharmacist divides a patient's supply of a pharmaceutical from a bulk supply, in that the patient always has access to the package insert contained in the patient pack, normally missing in traditional prescriptions. The inclusion of a package insert has been shown to improve patient compliance with the physician's instructions and, therefore, lead generally to more successful treatment. It will be understood that the administration of the combination of the invention by means of a single patient pack, or patient packs of each component compound, and containing a package insert instructing the patient to the correct use of the invention is a desirable additional feature of the invention.

Hereinafter, the term "active ingredients" means salmeterol or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, preferably salmeterol xinafoate, and mometasone, or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, preferably mometasone furoate.

Suitably, the pharmaceutical formulations which are suitable for inhalation according to the invention comprise the active ingredients in amounts such that each actuation provides therapeutically effective dose, for example, a dose of salmeterol of 10mcg to 150mcg, preferably 50mcg and a dose of mometasone of 100mcg to 1.6mg, preferably 200mcg to 1mg, more preferably, 200mcg to 400mcg.

The pharmaceutical formulations according to the invention may further include other therapeutic agents for example anti-inflammatory agents such as other corticosteroids (e.g. fluticasone propionate, beclomethasone dipropionate,

budenoside, or triamcinolone acetonide), or NSAIDs (e.g. sodium cromoglycate, nedocromil sodium, PDE-4 inhibitors, leukotriene antagonists, iNOS inhibitors, tryptase and elastase inhibitors, beta-2 integrin antagonists and adenosine 2a agonists), or other β_2 -adrenoreceptor agonists (such as salbutamol, formoterol, fenoterol or terbutaline and salts thereof), or anticholinergic agents (such as ipratropium, or tiotropium).

The formulations include those suitable for oral, parenteral (including subcutaneous, intradermal, intramuscular, intravenous and intraarticular), inhalation (including fine particle dusts or mists which may be generated by means of various types of metered dose pressurised aerosols, nebulisers or insufflators), rectal and topical (including dermal, buccal, sublingual and intraocular) administration although the most suitable route may depend upon for example the condition and disorder of the recipient. The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing the active ingredients into association with the carrier which constitutes one or more accessory ingredients. In general the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both and then, if necessary, shaping the product into the desired formulation.

Formulations for inhalation include powder compositions which will preferably contain lactose, and spray compositions which may be formulated, for example, as aqueous solutions or suspensions or as aerosols delivered from pressurised packs, with the use of a suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, 1,1,1,2,3,3,3-heptafluoropropane, 1,1,1,2-tetrafluoroethane, carbon dioxide or other suitable gas. Suitable aerosol formulations include those described in EP 0372777 and WO93/11743. For suspension aerosols, the active ingredients should be micronised so as to permit inhalation of substantially all of the active ingredients into the lungs upon administration of the aerosol formulation, thus the active ingredients will have a particle size of less than 100 microns, desirably less than 20 microns, and preferably in the range 1 to 10 microns, for example, 1 to 5 microns.

Intranasal sprays may be formulated with aqueous or non-aqueous vehicles with the addition of agents such as thickening agents, buffer salts or acid or alkali to adjust the pH, isotonicity adjusting agents or anti-oxidants.

5

Capsules and cartridges or for example gelatin, or blisters of for example laminated aluminium foil, for use in an inhaler or insufflator may be formulated containing a powder mix of the active ingredients and a suitable powder base such as lactose or starch. In this aspect, the active ingredients are suitably micronised so as to permit inhalation of substantially all of the active ingredients into the lungs upon administration of the dry powder formulation, thus the active ingredients will have a particle size of less than 100 microns, desirably less than 20 microns, and preferably in the range 1 to 10 microns.

10

15

Solutions for inhalation by nebulation may be formulated with an aqueous vehicle with the addition of agents such as acid or alkali, buffer salts, isotonicity adjusting agents or antimicrobials. They may be sterilised by filtration or heating in an autoclave, or presented as a non-sterile product.

20

Preferred unit dosage formulations are those containing a pharmaceutically effective dose, as hereinbefore recited, or an appropriate fraction thereof, of the active ingredient. Thus, in the case of formulations designed for delivery by metered dose pressurised aerosols, one actuation of the aerosol may deliver half of the therapeutically effective amount such that two actuations are necessary to deliver the therapeutically effective dose.

25

It should be understood that in addition to the ingredients particularly mentioned above, the formulations of this invention may include other agents conventional in the art having regard to the type of formulation in question. Furthermore, the claimed formulations include bioequivalents as defined by the US Food and Drugs Agency.

30

For a better understanding of the invention, the following Examples are given by way of illustration.

35

EXAMPLES**A: Metered Dose Inhalers****Example 1**

5

	Per actuation
Salmeterol Xinafoate	36.3 microgram
Mometasone	200 microgram
1,1,1,2-Tetrafluoroethane	to 75.0mg

The micronised active ingredients are weighed into an aluminium can, 1,1,1,2-tetrafluoroethane is then added from a vacuum flask and a metering valve is

10

Similar methods may be used for the formulation of Examples 2 and 3:

Example 2

15

	Per actuation
Salmeterol Xinafoate	36.3 microgram
Mometasone	100 microgram
1,1,1,2-Tetrafluoroethane	to 75.0mg

Example 3

	Per actuation
Salmeterol Xinafoate	36.3 microgram
Mometasone Furoate	100 microgram
1,1,1,2-Tetrafluoroethane	to 37.50mg

20

Example 4

	Per actuation
Salmeterol Xinafoate	36.3 microgram
Mometasone Furoate	100 microgram
1,1,1,2-Tetrafluoroethane	to 75.0mg

- 5 Salmeterol xinafoate (5.8mg) and mometasone furoate (16.0mg) were weighed directly into an 8ml aluminium canister coated internally with a PTFE/PES polymer blend as described in WO96/32150. A Valois DF60 metering valve was crimped into place, 1,1,1,2-tetrafluoroethane (to 12g) added and the filled canister was sonicated for at least five minutes. The resultant aerosol delivered
- 10 25 microgram salmeterol (as the xinafoate salt) and 100 microgram mometasone furoate per actuation.

An alternative method for preparing the formulations described in Examples 1 to 4 involves mixing the micronised medicaments and a portion of the propellant in a pressure vessel. An aliquot of the resultant suspension, followed by an aliquot of propellant is filled into a closed canister via the metering valve.

15

B: Dry Powder Inhalers

Example 5

20

	Per cartridge or blister
Salmeterol Xinafoate	72.6 microgram
Mometasone	200 microgram
Lactose Ph. Eur.	to 12.5mg or to 25.0mg

The active ingredients are micronised and bulk blended with the lactose in the proportions given above. The blend is filled into hard gelatin capsules or cartridges or in specifically constructed double foil blister packs to be

administered by an inhaler such as a Rotahaler, Diskhaler, or Diskus inhaler (each of these being a Trademark of Glaxo Group Limited).

Similar methods may be used for the formulations of Example 6:

5 Example 6

	Per cartridge or blister
Salmeterol Xinafoate	72.6 microgram
Mometasone	100 microgram
Lactose Ph. Eur.	to 12.5mg or to 25.0mg

Example 7

	Per blister
Salmeterol Xinafoate	72.5 microgram
Mometasone Furoate	200.0 microgram
Lactose Ph. Eur.	to 25.0mg

10

The active ingredients were micronised and bulk blended with the lactose in the proportions given above. The blend was then filled into specifically constructed double foil blister packs to be administered by a Diskhaler (Trademark of Glaxo Group Limited).

C: Suspension for nebulisation

Example 8

	Quantity (mg)
Salmeterol Xinafoate (micronised)	0.0725
Mometasone Furoate (micronised)	0.20
Polysorbate 20	0.14
Sorbitan Monolaurate	0.018
Monosodium Phosphate dihydrate	18.80
Dibasic Sodium Phosphate anhydrous	3.50
Sodium Chloride	9.60
Water for injection	to 2.00 ml

5

D: Aqueous nasal spray

Example 9

	Quantity ¹ (%w/w)
Salmeterol Xinafoate (micronised)	0.03625
Mometasone Furoate (micronised)	0.05
Dextrose Anhydrous	5.00
Microcrystalline cellulose and carboxymethylcellulose sodium	1.50
Phenylethyl alcohol	0.25
Benzalkonium Chloride solution (50%w/v)	0.04v/w
Polysorbate 80	0.005
Purified water	to 100

¹ Based on 100mg suspension per actuation

10

E: Intranasal dry powder

Example 10

	Per blister
Salmeterol Xinafoate (micronised)	72.50 microgram
Mometasone Furoate (micronised)	100.00 microgram
Potato Starch NF/BP	to 10mg

5 Cascade Impaction Data

The particle size distribution of the aerosol formulations according to the invention may be measured by conventional techniques, such as cascade impaction (for example as defined in US Pharmacopoeia, 23/NF18 General Test <601>, pages 1762 - 1765).

Results

For the product of Example 4, the Cascade Impaction data were as follows:

15 Salmeterol:

Sample	Sample 1 Total %	Sample 2 Total %	Mean Total %
Device	12.1	15.6	13.8
Throat	35.2	36.5	35.8
Stage 0	6.5	4.1	5.3
Stage 1	2.4	2.0	2.4
Stage 2	2.8	2.5	2.8
Stage 3	13.0	11.5	12.2
Stage 4	15.8	15.6	15.9
Stage 5	11.3	11.5	11.4
Stage 6	0.8	0.8	0.8
Stage 7	0.0	0.0	0.0
Filter	0.0	0.0	0.0
Total	100.0	100.0	100.0

13

Sample	Sample 1 Total %	Sample 2 Total %	Mean Total %
Total Ex-device	87.9	84.4	86.2
FPM Sum Stages 3-5	40.1	38.5	39.4

Mometasone:

Sample	Sample 1 Total %	Sample 2 Total %	Mean Total %
Device	6.0	9.2	7.7
Throat	34.0	38.0	36.0
Stage 0	6.2	3.9	5.0
Stage 1	2.6	2.4	2.5
Stage 2	3.0	2.7	2.8
Stage 3	19.5	16.3	17.9
Stage 4	18.6	17.2	17.9
Stage 5	9.3	9.3	9.3
Stage 6	0.7	0.6	0.7
Stage 7	0.2	0.2	0.2
Filter	0.0	0.2	0.1
Total	100.0	100.0	100.0
Total Ex-device	94.0	90.8	92.3
FPM Sum Stages 3-5	47.3	42.8	45.0

5 These data are also presented graphically in Figure 1.

For the product of Example 7, the Cascade Impaction data were as follows

Salmeterol:

Sample	Disk 2 Total %	Disk 4 Total %	Mean Total %
Throat	16.4	16.0	16.3
Pre-sep & Stage 0	51.9	50.6	51.2
Stage 1	5.0	5.5	5.2
Stage 2	3.9	4.0	3.9
Stage 3	10.6	11.2	11.0
Stage 4	8.2	8.1	8.0
Stage 5	2.8	3.5	3.1
Stage 6	0.8	0.7	0.8
Stage 7	0.6	0.2	0.5
Filter	0.0	0.0	0.0
Total	100.0	100.0	100.0
FPM Sum Stages 1-5	30.3	32.4	31.5

Mometasone:

5

Sample	Disk 2 Total %	Disk 4 Total %	Mean Total %
Throat	15.2	15.0	15.1
Pre-sep & Stage 0	48.7	46.2	47.4
Stage 1	6.4	6.7	6.5
Stage 2	5.5	6.0	5.8
Stage 3	13.2	14.6	13.9
Stage 4	7.1	7.5	7.3
Stage 5	2.6	3.1	2.9
Stage 6	0.8	0.8	0.8
Stage 7	0.5	0.3	0.4
Filter	0.0	0.0	0.0
Total	100.0	100.0	100.0
FPM Sum	34.8	37.8	36.4

15

Sample	Disk 2 Total %	Disk 4 Total %	Mean Total %
Stages 1-5			

These data are also presented graphically in Figure 2.

Claims

1. A pharmaceutical formulation comprising salmeterol or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof and mometasone or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, and a pharmaceutically acceptable carrier or excipient, and optionally one or more other therapeutic ingredients.
2. A pharmaceutical formulation comprising salmeterol xinafoate and mometasone furoate, and a pharmaceutically acceptable carrier or excipient, and optionally one or more other therapeutic ingredients.
3. A pharmaceutical formulation according to claim 1 or 2 which is suitable for administration by inhalation.
4. A pharmaceutical formulation according to any of claims 1 to 3 wherein the pharmaceutically acceptable carrier or excipient is lactose.
5. A pharmaceutical formulation according to any of claims 1 to 3 wherein the pharmaceutically acceptable carrier or excipient comprises 1,1,1,2-tetrafluoroethane and/or 1,1,1,2,3,3,3- heptafluoropropane.
6. A method for the prophylaxis or treatment of a clinical condition in a mammal, such as a human, for which a selective β_2 -adrenoreceptor agonist and/or antiinflammatory corticosteroid is indicated, which comprises administration of a therapeutically effective amount of a pharmaceutical formulation according to any one of claims 1 to 5.
7. A method according to claim 6 wherein the clinical condition is a disease associated with reversible airways obstruction such as asthma, chronic obstructive pulmonary disease (COPD), respiratory tract infection or upper respiratory tract disease.





FIGURE 2
Cascade Impaction Profile
Mean % Results



INTERNATIONAL SEARCH REPORT

Int'l Application No

PCT/GB 01/01637

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/575 A61K31/137 A61P11/06 //(A61K31/575,31:137)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, MEDLINE, BIOSIS, CHEM ABS Data, EMBASE, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 98 41193 A (SCHERING CORP) 24 September 1998 (1998-09-24) cited in the application claims 1,5,46-50	1-7
Y	US 4 472 393 A (SHAPIRO ELLIOT L) 18 September 1984 (1984-09-18) cited in the application the whole document	1-7
	-/-	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

8 August 2001

Date of mailing of the international search report

04/09/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Herrera, S

INTERNATIONAL SEARCH REPORT

Int. Application No

PCT/GB 01/01637

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	BARNES P J ET AL: "EFFICACY OF INHALED CORTICOSTEROIDS IN ASTHMA" JOURNAL OF ALLERGY AND CLINICAL IMMUNOLOGY, MOSBY - YEARLY BOOK, INC, US, vol. 102, no. 4, 1998, pages 531-538, XP000913470 ISSN: 0091-6749 page 536, right-hand column, line 1-6 abstract	1-7
Y	O'CONNOR B J: "COMBINATION THERAPY" PULMONARY PHARMACOLOGY AND THERAPEUTICS, ACADEMIC PRESS, NEW YORK, NY, US, vol. 11, no. 5/6, 1998, pages 397-399, XP000911059 ISSN: 1094-5539 the whole document	1-7
Y	BOWLER S: "LONG ACTING BETA AGONISTS" AUSTRALIAN FAMILY PHYSICIAN, XX, XX, vol. 27, no. 12, December 1998 (1998-12), pages 1115,1117-1118, XP000973076 the whole document	1-7
Y	GB 2 140 800 A (GLAXO GROUP LTD) 5 December 1984 (1984-12-05) cited in the application the whole document	1-7
Y	WILDING PAUL ET AL: "Effect of long term treatment with salmeterol on asthma control: A double blind, randomised crossover study." BMJ, vol. 314, no. 7092, 1997, pages 1441-1446, XP001013219 page 1445, right-hand column, line 1-15	1-7

INTERNATIONAL SEARCH REPORT

Inte al Application No

PCT/GB 01/01637

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9841193 A	24-09-1998	AU 6537898 A	12-10-1998
		CN 1257423 T	21-06-2000
		EP 0969816 A	12-01-2000
		HU 0002029 A	28-11-2000
		JP 2000510478 T	15-08-2000
		NO 994519 A	19-11-1999
		PL 335742 A	08-05-2000
		SK 128099 A	12-06-2000
		ZA 9802254 A	17-09-1998
US 4472393 A	18-09-1984	AT 8790 T	15-08-1984
		AU 549102 B	16-01-1986
		AU 7991882 A	12-08-1982
		BG 60799 B	29-03-1996
		CA 1177822 A	13-11-1984
		CY 1359 A	07-08-1987
		DE 3260474 D	06-09-1984
		DK 39082 A,B,	03-08-1982
		EP 0057401 A	11-08-1982
		FI 820280 A,B,	03-08-1982
		HK 68487 A	02-10-1987
		HU 188769 B	28-05-1986
		IE 52576 B	23-12-1987
		IL 64885 A	31-05-1985
		JP 1512102 C	09-08-1989
		JP 57146800 A	10-09-1982
		JP 63060036 B	22-11-1988
		KE 3694 A	13-03-1987
		KR 8900761 B	05-04-1989
		MX 9203403 A	01-07-1992
		NO 820263 A,B,	03-08-1982
		NZ 199600 A	28-09-1984
		OA 7116 A	31-03-1984
		PH 19733 A	17-06-1986
		PT 74357 A,B	01-02-1982
		ZA 8200566 A	29-12-1982
GB 2140800 A	05-12-1984	AT 390611 B	11-06-1990
		AT 129184 A	15-11-1989
		AU 573212 B	02-06-1988
		AU 2706484 A	25-10-1984
		BE 899448 A	18-10-1984
		BE 900835 A	17-04-1985
		BG 61490 B	30-09-1997
		CA 1335999 A	20-06-1995
		CA 1336004 A	20-06-1995
		CH 661497 A	31-07-1987
		CH 667084 A	15-09-1988
		CZ 9104028 A	12-05-1993
		CY 1482 A	08-12-1989
		CZ 285602 B	15-09-1999
		DE 3414752 A	18-10-1984
		DE 3448338 C	19-05-1993
		DE 3448452 C	05-01-1994
		DK 158092 A	30-12-1992
		DK 201784 A	19-10-1984
		ES 531722 D	01-06-1985
		ES 8505641 A	01-10-1985

INTERNATIONAL SEARCH REPORT

formation on patent family members

International Application No

PCT/GB 01/01637

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 2140800 A		ES 539625 D	16-07-1986
		ES 8609209 A	16-12-1986
		FI 841548 A,B,	19-10-1984
		FR 2545482 A	09-11-1984
		GB 2176476 A,B	31-12-1986
		GR 79925 A	31-10-1984
		HK 36889 A	12-05-1989
		HU 200160 B	28-04-1990
		IE 57237 B	17-06-1992
		IL 71569 A	30-10-1987
		IT 1199112 B	30-12-1988
		JP 1862136 C	08-08-1994
		JP 5069817 B	01-10-1993
		JP 63264443 A	01-11-1988
		JP 1643826 C	28-02-1992
		JP 3000858 B	09-01-1991
		JP 59199659 A	12-11-1984
		JP 2001385 C	20-12-1995
		JP 6087800 A	29-03-1994
		JP 7029997 B	05-04-1995
		KE 3864 A	19-05-1989
		KR 9204186 B	30-05-1992
		LU 85329 A	04-06-1985
		LU 88265 A	03-02-1994
		MX 9203226 A	01-07-1992
		NL 8401258 A,B,	16-11-1984
		NO 841568 A,B,	19-10-1984
		NZ 207885 A	29-11-1988
		NZ 221999 A	29-11-1988